

2

AD-A204 970

REPORT DOCUMENTATION PAGE

1b. RESTRICTIVE MARKINGS	
3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NMRI 88-77	
5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Naval Medical Research	6b. OFFICE SYMBOL (if applicable)
7a. NAME OF MONITORING ORGANIZATION Naval Medical Command	
7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Washington, D.C. 20372-5120	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Naval Medical Research and Development Command	8b. OFFICE SYMBOL (if applicable)
9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
10. SOURCE OF FUNDING NUMBERS	
PROGRAM ELEMENT NO. 61153N	PROJECT NO. MR04101
TASK NO. 01.1005	WORK UNIT ACCESSION NO. DN277001
11. TITLE (Include Security Classification) Pressure suppresses serotonin release by guinea pig striatal synaptosomes	
12. PERSONAL AUTHOR(S) Gilman SC, Colton JS, Hsu SC, Dutka AJ	
13a. TYPE OF REPORT journal article	13b. TIME COVERED FROM _____ TO _____
14. DATE OF REPORT (Year, Month, Day) 1988	
15. PAGE COUNT 9	
16. SUPPLEMENTARY NOTATION Reprinted from: Undersea Biomedical Research 1988 Vol.15(2) pp. 69-77	
17. COSATI CODES	
FIELD	GROUP
SUB-GROUP	
18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Serotonin efflux High-pressure nervous syndrome Synaptosome Striatum	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)	
<p style="text-align: center;">DTIC ELECTED 24 FEB 1989 E</p>	
Accession For	
NTIS GRA&I <input checked="" type="checkbox"/>	
DTIC TAB <input checked="" type="checkbox"/>	
Unannounced <input type="checkbox"/>	
Justification <input type="checkbox"/>	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-120	
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS	
21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Phyllis Blum, Information Services Division	
22b. TELEPHONE (Include Area Code) 202-295-2188	
22c. OFFICE SYMBOL ISD/ADMIN/NMRI	

Pressure suppresses serotonin release by guinea pig striatal synaptosomes

S.C. GILMAN, J.S. COLTON, S.C. HSU, and A.J. DUTKA

Diving Medicine Department, Naval Medical Research Institute, Bethesda, MD 20814-5055

embolism in

^{+ tritium}
Gilman SC, Colton JS, Hsu SC, Dutka AJ. Pressure suppresses serotonin release by guinea pig striatal synaptosomes. Undersea Biomed Res 1988; 15(2):69-77—Exposure to high pressure produces neurologic changes in humans which manifest as tremor, EEG changes, and convulsions. Since previous studies have implicated the involvement of the serotonergic system in these symptoms, it was of interest to study serotonin release at high pressure. Synaptosomes isolated from guinea pig striatum were used to follow serotonin efflux at 68 ATA. The major observation was a decrease in ³H serotonin release from depolarized striatal synaptosomes at 68 ATA. In view of the role of serotonin as an inhibitory neurotransmitter in this area, the observed decrease in synaptic release leads us to conclude that decreased serotonergic activity in striatal neurons probably is contributing to the hyperexcitability associated with HPNS.

Stress Physiology Reprints (AW)
serotonin efflux synaptosome
high pressure nervous syndrome striatum

INTRODUCTION

→ ^{start} Exposure to pressures greater than 27 atmospheres absolute (ATA) produces a disturbance of neurologic function referred to as the high pressure neurologic syndrome (HPNS) (1). Neurologic changes of HPNS manifest as tremor, myoclonic episodes, EEG changes, and convulsions (2, 3).

Although a number of pharmacologic agents have been tested to protect against the adverse effects of pressure (4, 5), little information is available about the direct action of high pressure on synaptic transmission in the CNS. Gillard et al. (6) recently used a differential pulse voltammetry technique to study the generation of serotonin metabolites in the rat striatum during exposures to high pressure. They found that exposure to 61-121 ATA produced a significant increase in the striatal concentration of 5-HIAA, the major metabolite of serotonin.

This finding suggests that pressure accelerates the turnover of serotonin in presynaptic nerve terminals in the striatum. Such an alteration in receptor function would be expected to affect the release kinetics of the striatal serotonergic system. There-

fore, we used an in vitro presynaptic nerve terminal preparation to determine the extent to which the synaptic release of [^3H]serotonin by striatal neurons is affected by high pressure exposure. The synaptic vesicle (synaptosome) preparation was chosen because it has been shown to retain the transport properties of the nerve ending in situ (7-9) and to exhibit a calcium-dependent release of neurotransmitter substances (10). Also, synaptosomes offered an advantage as an experimental model in that multicellular or neuroglial interactions could be eliminated.

MATERIAL AND METHODS

Animals

Adult male Hartley short-haired guinea pigs (300 and 400 g) were obtained from a commercial source.

Materials

Radioactive 5-[1,2- $^3\text{H}(\text{N})$]hydroxytryptamine binoxalate (serotonin; 15-30 Ci/mmol) was purchased from New England Nuclear Corporation, Boston, MA. All other chemicals were purchased from Sigma Chemical Company, St. Louis, MO.

Synaptosomal isolation procedure

Each guinea pig was killed by decapitation, and the striatum dissected into 10 vol of 0.32 M sucrose buffered with 5 mM HEPES. Synaptosomes were prepared by density gradient centrifugation as previously described (11). Briefly, a homogenizer was used to disperse the tissue. Centrifugations were carried out at 3°C in an ultracentrifuge using a digital integrator to obtain consistent centrifugation. The initial homogenate was centrifuged at $1100 \times g$ for 5 min to yield a nuclear pellet and low speed supernatant. The supernatant was centrifuged at $10,000 \times g$ for 10 min to yield a mitochondrial pellet which was resuspended in sucrose and homogenized. This homogenate was layered over a Ficoll gradient and centrifuged at $26,000 \times g$ for 30 min. After centrifugation, the purified synaptosomes were removed, diluted with sucrose, and pelleted.

Radioisotope uptake

The method used for [^3H]serotonin loading and efflux measurement of synaptosomes was developed by modification of the methods of Gilman (12) and Schlicker et al. (13). The final synaptosome pellet was suspended in 20 vol of buffer. [^3H]Serotonin was added and the mixture was incubated at 37°C for 30 min to allow uptake by the tissue. After uptake, a 1.0-ml aliquot of the synaptosome suspension was placed on a filter unit consisting of a 0.45- μm nylon membrane filter positioned on the plastic multiperforated support of a 10-ml perfusion chamber. The perfusion chamber was then connected to a polystaltic pump, and the filters immediately washed with 25 ml of a high sodium, calcium-free buffer medium at 37°C, using the highest pump speed.

Compression studies

After the [^3H]serotonin-loaded synaptosomes were washed, the filter unit was attached to a superfusion apparatus that was mounted in a hyperbaric chamber. The hyperbaric chamber was sealed and pressurized to 1.3 ATA with oxygen, and to a final pressure of 68 ATA with helium. The rate of compression was 4 ATA/min. Oxygen partial pressure, as measured with a paramagnetic O_2 analyzer, was maintained at 0.49 ± 0.01 ATA throughout pressurization. A recirculation atmosphere control system was used to regulate CO_2 at < 0.0005 ATA, as measured by an infrared analyzer. Temperature within the hyperbaric chamber was maintained at $37^\circ \pm 2^\circ\text{C}$ during compression, as previously described (13). During pressurization and the release studies the temperatures of the buffer media and perfusion chamber were constantly monitored by microthermistors, and maintained at $37^\circ \pm 0.5^\circ\text{C}$ using YSI model 73A temperature control units (Yellow Springs Instrument Co., Yellow Springs, OH).

Upon reaching 68 ATA, the filters were perfused at a rate of 500 $\mu\text{l}/\text{min}$ with 5 ml of buffer medium and 3 perfusate "wash" fractions collected directly into scintillation vials. At 3 min after the beginning of superfusion, the buffer medium was quickly replaced with 10 ml of a 5 mM K^+ , nondepolarizing or 10 ml of a depolarizing, 70 mM K^+ "efflux" medium containing 1.2 ml of Ca^{2+} . The composition was (in millimolars): NaCl, 145 or 80; KCl, 5 or 70; CaCl_2 , 2.5; MgCl_2 , 1.2; KH_2PO_4 , 1.2; HEPES, 20; glucose, 10; pH 7.4. Nine fractions containing 500 μl each of perfusate were collected every minute directly into scintillation vials. The hyperbaric chamber was then decompressed to 1 ATA at the rate of 4.03 ATA/min. In all experiments, aliquots from the same synaptosome preparation were used for obtaining the 1 and 68 ATA release values. Control aliquots from the same preparation were treated identically to the experimental group, except that pressure exposures were sham (1 ATA).

Assay of samples

Biofluor (15 ml) was added to the scintillation vial and the radioactivity of each perfusate was determined. The radioactivity remaining on the filters at the end of the superfusion was also counted. Each filter was placed in a scintillation vial containing 500 μl of 1% sodium dodecyl sulfate. After agitation, 15 μl of Biofluor was added and the radioactivity of the filter was measured.

Expression of results

Fractional efflux was expressed as percentage released per minute of total radioactivity, i.e.,

$$\text{efflux} = \frac{\text{cpm in 500 } \mu\text{l filtrate}}{\text{total radioactivity}}$$

where total radioactivity was the sum of all 12 fractional filtrate cpm's and cpm remaining on the filter. Statistical significances at each time point were determined using *t* tests.

RESULTS

 $[^3\text{H}]$ Serotonin release at 1 ATA

Figure 1 shows the time course of depolarization-induced efflux of $[^3\text{H}]$ serotonin from synaptosome fractions isolated from the striatum. Application of the high K^+ medium containing Ca^{2+} initially induced a more than fourfold increase in the release of $[^3\text{H}]$ serotonin by the synaptosome preparation. After the first 3 min, the evoked release declined rapidly, returning within 8 min to levels not significantly greater than the resting efflux levels.

Synaptosomes were perfused with a depolarizing, 70 mM K^+ calcium-free medium, with MgCl_2 replacing the omitted CaCl_2 to determine the extent to which the release observed with depolarization was calcium-dependent. Figure 1 shows the effect of calcium removal on depolarization-induced $[^3\text{H}]$ serotonin release. As can be seen, the removal of calcium from the depolarizing medium reduced initial release evoked by high K^+ by over 80%.

Compression effects on $[^3\text{H}]$ serotonin

Figure 2 shows the effect of compression to 68 ATA on the nondepolarized release of $[^3\text{H}]$ serotonin. Pressure was not found to cause any significant changes in the spontaneous efflux of serotonin by these striatal synaptosomes ($P > 0.05$).

As shown in Fig. 3 A, the evoked release of $[^3\text{H}]$ serotonin from synaptosomes in the presence of 1.2 mM Ca^{2+} was substantially reduced (i.e., 65%) by compression




Fig. 1. Release of $[^3\text{H}]$ serotonin from guinea pig striatal synaptosome fractions. *Triangle*, depolarized, Ca^{2+} present; *circle*, depolarized, Ca^{2+} absent; *square*, nondepolarized, Ca^{2+} present. After preloading with $[^3\text{H}]$ serotonin, synaptosome aliquots were superfused with a high sodium, calcium-free buffer medium for 3 min, then 70-mM K^+ -containing medium with or without Ca^{2+} or 5-mM K^+ -containing medium with Ca^{2+} was similarly superfused for the 9 min marked by horizontal bar (i.e., 3–12 min).




Fig. 2. Effect of compression to 68 ATA on the spontaneous (5-mM K^+ -evoked) release of [3H]serotonin. Triangle, 1 ATA; circle, 68 ATA.

to 68 ATA. Analysis of the data showed that the depression in serotonin release by these synaptic terminals was significant at the 4-min superfusion time ($P < 0.05$). On the other hand, the release in the absence of Ca^{2+} was little affected by compression ($P > 0.05$; Fig. 3 B). Figure 3 C shows the difference in [3H]serotonin release between the Ca^{2+} -containing and Ca^{2+} -free media, which represents the calcium-dependent component of [3H]serotonin release. Statistical comparison between the 1 ATA control curve and the 68 ATA calcium-dependent curve showed that [3H]serotonin release at the 4-min superfusion time was significantly lower in the presence of pressure than in the corresponding control condition ($P < 0.05$). This indicates that pressure exposure depressed the initial depolarization-induced, calcium-dependent release of [3H]serotonin by striatal synaptosomes isolated from the striatum.

DISCUSSION

Electron microscopic examination and the measurement of the cytoplasmic markers showed a high purity synaptosome preparation which retains an intact plasma membrane, with no apparent nonselective changes in permeability, which would be reflective of membrane damage. Also important to data interpretation was that high pressure exposure has not been found to have any disruptive effect on the synaptosomal plasma membrane (13).

High K^+ medium containing 1.2 mM Ca^{2+} initially produced a rapid increase in the release of [3H]serotonin by the synaptosome preparation, followed by a decline in release approaching resting efflux levels. The removal of calcium from the depolarizing medium reduced [3H]serotonin release evoked by high potassium by over 80%. This indicates that the evoked release of serotonin by striatal synaptosomes is highly dependent on the influx of calcium.




Fig. 3. Effect of compression to 68 ATA on the high K^+ -evoked release of $[^3H]$ serotonin from striatal synaptosomes. A, in the presence of Ca^{2+} ; B, in the absence of Ca^{2+} ; C, Ca^{2+} -dependent release, i.e., the difference between A and B. Triangle, 1 ATA; circle, 67.7 ATA. $\star p < .05$.

The major observation in the present study was a 65% decrease in the initial efflux of [^3H]serotonin from depolarized striatal synaptosomes after compression to 68 ATA. This suppression of serotonin release parallels decreases noted previously in the calcium-dependent release of other neurotransmitters with pressure (11, 14, 15).

The pressure-induced decrease in the initial rate (viz in the first 1–2 min) of transmitter efflux seen in this and other studies (11, 14, 15) suggests that a concomitant decrease in voltage-sensitive calcium influx might be occurring at high pressure. However, in another study (16) we have observed that although $^{45}\text{Ca}^{2+}$ movement into synaptosomes is significantly depressed at 68 ATA, the initial (i.e., the first minute) amount of Ca^{2+} influx, although depressed, was not significantly different from control values (1 ATA). This relationship between the initial amount of $^{45}\text{Ca}^{2+}$ uptake and the significant depression of initial transmitter efflux leads us to propose that high pressure helium may alter transmitter release in more than one way.

First, high pressure most likely does interfere with voltage-sensitive calcium channel gating. Second, high pressure also probably interrupts some aspect of the intraterminal cascade for the calcium-dependent release of transmitter (e.g., calcium-calmodulin interaction, kinase II activity, membrane phosphorylation, or the interaction of the vesicle membrane with the terminal membrane) (16–18). Further support for this notion is provided by a preliminary study in our laboratory with the calcium ionophore, A23187, applied to synaptosomes subjected to high pressure. This study indicates that this ionophore does not alter the decrease in transmitter release at 68 ATA. A23187 circumvents the membrane-bound Ca^{2+} channel by forming a Ca^{2+} - H^{+} exchanger and therefore should facilitate the rise in intraterminal Ca^{2+} (19, 20). Since the effect of high pressure appears to be universal on calcium-dependent transmitter release, alteration of any or all of the above-mentioned sites could affect transmitter release at most synapses.

The increase in the serotonin metabolite, 5-HIAA, found by Gillard et al. (6) suggests an increase in metabolic activity or a decrease in the removal of this compound. Removal of 5-HIAA from the brain requires a membrane-bound protein carrier. Inhibition of this carrier by the drug Probenecid has been shown to produce an accumulation of 5-HIAA in brain tissue (21, 22). Since removal requires a membrane-bound protein carrier, it is reasonable to suspect that high pressure helium could exert its effect on the membrane to depress carrier activity, causing 5-HIAA levels to rise. It is unlikely that this inhibition of metabolite removal suppresses serotonin production, because serotonin synthesis does not appear to be feedback regulated (23). Therefore, the depression of transmitter release found in our study is not adequately explained by changes in serotonin metabolism or metabolite removal.

Whether this pressure-induced phenomenon is a contributory factor in the actual development of the symptoms of HPNS is currently unknown. However, the decrease in the initial rate of release of serotonin observed in synaptosomes may reflect in vivo events occurring in the synaptic terminal.

Since serotonin is believed to function as a modulator to presynaptically inhibit transmitter release (24, 25), a small change in the timing of release of serotonin could result in a marked diminution of its inhibitory function. This could result in a significantly enhanced excitatory response. Coupled with similar events occurring in other inhibitory systems (viz GABA and glycine), symptomatic neurologic changes characteristic of HPNS could result.

The authors give special thanks to Jeff Boogaard and Alex Lee for valuable technical support. The authors also thank Susan Cecire for her assistance in the preparation of this manuscript.

This study was supported by the Naval Medical Research and Development Command Research Task MR041.01.01.1124.

The opinions and assertions contained herein are the private ones of the writers and are not to be construed as official or reflecting the views of the Navy Department of the naval service at large. The experiments reported herein were conducted according to the principles set forth in the current edition of the "Guide for the Care and Use of Laboratory Animals," Institute of Laboratory Animal Resources, National Research Council.—*Manuscript received March 1987; accepted December 1987.*

Gilman SC, Colton JS, Hsu SC, Dutka AJ. La pression supprime la libération de sérotonine par les synaptosomes striataires de cobaye.—*Undersea Biomed Res* 1988; 15(2):69–77. L'exposition à la pression élevée produit des changements neurologiques chez les humains, lesquels se manifestent par le tremblement, des changements dans l'EEG et des convulsions. Puisque des études antérieures ont impliqué le système nerveux sérotoninergique dans ces symptômes, il est d'intérêt à étudier la libération de sérotonine sous haute pression. Des synaptosomes isolés du corps strié de cobaye furent utilisés pour suivre l'efflux de sérotonine à 68 ATA. L'observation principale fut une diminution dans la libération de [³H]sérotonine de synaptosomes striataires dépolarisés à 68 ATA. Vu le rôle de la sérotonine comme neurotransmetteur inhibiteur dans cette zone, la diminution observée dans la libération synaptique mène à conclure que la réduction de l'activité sérotoninergique dans les neurones striataires contribue probablement à la surexcitabilité associée avec le syndrome nerveux de la haute pression.

Gilman SC, Colton JS, Hsu SC, Dutka AJ. Supresión por presión de la liberación de serotonina en sinaptosomas del cuerpo estriado de conejillos de Indias.—*Undersea Biomed Res* 1988; 15(2):69–77. La exposición a presiones altas produce cambios neurológicos en humanos, que se manifiestan como temblor, cambios electroencefalográficos y convulsiones. Debido a que trabajos previos han implicado la participación del sistema serotoninérgico en estos síntomas, nuestro interés residió en el estudio de la liberación de serotonina a presiones altas. Se empleó sinaptosomas del cuerpo estriado de conejillos de indias para estudiar el flujo de serotonina a 68 ATA. El hallazgo más sobresaliente fue la disminución en la liberación de [³H]serotonina en sinaptosomas despolarizados de cuerpo estriado a 68 ATA. Debido a que la función de la serotonina a este nivel es de un neurotransmisor inhibitorio, la disminución observada en la liberación sináptica nos lleva a concluir que la reducción en la actividad serotoninérgica en las neuronas del cuerpo estriado probablemente contribuye a la hiperexcitabilidad asociada al Síndrome Neurológico de Presión Alta (SNPA).

REFERENCES

1. Brauer RW. The high pressure nervous syndrome: animals. In: Bennett PB, Elliot DH, eds. *The physiology and medicine of diving and compressed air work*. London: Ballière-Tindall, 1975:230–247.
2. Hallenbeck JM. A view of some fundaments of the high pressure nervous syndrome. In: Bachrach AJ, Matzen MM, eds. *Undersea physiology VII. Proceedings of the seventh symposium on underwater physiology*. Bethesda, MD: Undersea Medical Society, 1981:305–315.
3. Hunter WL, Bennett PB. The causes, mechanisms, and prevention of the high pressure nervous syndrome. *Undersea Biomed Res* 1974; 1:1–28.
4. Bichard AR, Little HJ. Drugs that increase γ -aminobutyric acid transmission protect against the high pressure neurological syndrome. *Brit J Pharmacol* 1982; 76:447–452.
5. Wardley-Smith B, Meldrum BS. Effect of excitatory amino acid antagonists on the high pressure neurological syndrome in rats. *Eur J Pharmacol* 1984; 105:351–354.

6. Gillard J, Bugat R, Obrenovitch J. Effects of high pressure of heliox on the striatal 5-HIAA and ascorbic acid rates in the rat. *Cent Etud Rech Bio-Physiol Rep* 84-08, 1984:35.
7. Blaustein MP, Kendrick NC, Fried RC, Ratzlaff RW. Calcium metabolism at the mammalian presynaptic nerve terminal: lessons from the synaptosome. In: Cowan WM, Ferrendelli, JA, eds. *Approaches to the cell biology of neurons*. Bethesda, MD: Society for Neuroscience, 1977:172-194.
8. Debellerroche JS, Bradford HF. The synaptosome: an isolated, working neuronal compartment. *Prog Neurobiol* 1973; 1:277-298.
9. Levy WB, Redburn DA, Cotman CW. Stimulus-coupled secretion of λ -aminobutyric acid from rat brain synaptosomes. *Science* 1973; 181:676-678.
10. Cotman CW, Haycock JW, White WF. Stimulus-secretion coupling processes in brain: analysis of noradrenaline and gamma-aminobutyric acid release. *J Physiol* 1976; 254:475-505.
11. Gilman SC, Kumaroo KK, Hallenbeck JM. Effect of pressure on the uptake and release of calcium by brain synaptosomes. *J Appl Physiol* 1986; 60:1446-1450.
12. Gilman SC. Effect of high pressure on calcium activity by guinea pig brain synaptosomes. Washington, DC: The George Washington University; 1985. Dissertation.
13. Schlicker E, Brandt F, Classen K, Bothert M. Serotonin release in human cerebral cortex and its modulation via serotonin receptors. *Brain Res* 1985; 332:337-341.
14. Gilman SC, Colton JS, Hallenbeck JM. Effect of pressure on [3 H]GABA release by synaptosomes isolated from the cerebral cortex. *J Appl Physiol* 1986; 61:2067-2073.
15. Gilman SC, Colton JS, Dutka AJ, Boogaard JS. Effect of high pressure on the release of excitatory amino acids by brain synaptosomes. *Undersea Biomed Res* 1986; 13:397-406.
16. Augustine GJ, Charlton MP, Smith SJ. Calcium action in synaptic transmitter release. *Ann Rev Neurosci* 1987; 10:633-693.
17. Huttner WB, Schiebler W, Greengard P, DeCamilli P. Synapsin I (protein I), a nerve terminal-specific phosphoprotein. III. Its association with synaptic vesicles studied in a highly purified synaptic vesicle preparation. *J Cell Biol* 1983; 96:1374-1388.
18. Llinas R, McGuinness TL, Leonard CS, Sugimori M, Greengard P. Intraterminal injection of synapsin I or calcium/calmodulin-dependent protein kinase II alters neurotransmitters release at squid giant synapse. *Proc Natl Acad Sci USA* 1985; 82:3035-3039.
19. Akermam KEO, Nicholls DG. Calcium transport by intact synaptosomes. Influence of ionophore A23187 on plasma-membrane potential, plasma membrane calcium transport, mitochondrial membrane potential, respiration, cytosolic free calcium concentration and noradrenaline release. *Eur J Biochem* 1981; 115:67-73.
20. Reed PW, Lardy HA. A23187: A divalent cation ionophore. *J Biol Chem* 1972; 247:6970-6977.
21. Neff NH, Tozer TN, Brodie BB. Application of steady-state kinetics to studies of the transfer of 5-hydroxyindoleacetic acid from brain to plasma. *J Pharmacol Exp Ther* 1967; 158:214-218.
22. Van Der Poel, FW, Van Proag HM, Kork J. Evidence for a probenecid-sensitive transport system of acid monoamine metabolites from the spinal subarachnoid space. *Psychopharmacology* 1977; 52:35-40.
23. Brownstein MJ. Serotonin, histamine and the purines. In: Siegel GJ, Albers RW, Agranoff BW, Katzman R, eds. *Basic neurochemistry*, third edition. Boston: Little Brown and Co. 1981:219-231.
24. Bennett JP, Solomon SH. Serotonin synaptic receptors in the mammalian central nervous system. *Fed Proc* 1978; 37:137-138.
25. Haigler HJ, Aghajanian GK. Serotonin receptors in the brain. *Fed Proc* 1977; 36:2159-2164.

6. Gillard J, Bugat R, Obrenovitch J. Effects of high pressure of heliox on the striatal 5-HIAA and ascorbic acid rates in the rat. *Cert Etud Rech Bio-Physiol Rep* 84-08, 1984:35.
7. Blaustein MP, Kendrick NC, Fried RC, Ratzlaff RW. Calcium metabolism at the mammalian presynaptic nerve terminal: lessons from the synaptosome. In: Cowan WM, Ferrendelli JA, eds. *Approaches to the cell biology of neurons*. Bethesda, MD: Society for Neuroscience, 1977:172-194.
8. Debellerocche JS, Bradford HF. The synaptosome: an isolated, working neuronal compartment. *Prog Neurobiol* 1973; 1:277-298.
9. Levy WB, Redburn DA, Cotman CW. Stimulus-coupled secretion of λ -aminobutyric acid from rat brain synaptosomes. *Science* 1973; 181:676-678.
10. Cotman CW, Haycock JW, White WF. Stimulus-secretion coupling processes in brain: analysis of noradrenaline and gamma-aminobutyric acid release. *J Physiol* 1976; 254:475-505.
11. Gilman SC, Kumaroo KK, Hallenbeck JM. Effect of pressure on the uptake and release of calcium by brain synaptosomes. *J Appl Physiol* 1986; 60:1446-1450.
12. Gilman SC. Effect of high pressure on calcium activity by guinea pig brain synaptosomes. Washington, DC: The George Washington University; 1985. Dissertation.
13. Schlicker E, Brandt F, Classen K, Bothert M. Serotonin release in human cerebral cortex and its modulation via serotonin receptors. *Brain Res* 1985; 332:337-341.
14. Gilman SC, Colton JS, Hallenbeck JM. Effect of pressure on [3 H]GABA release by synaptosomes isolated from the cerebral cortex. *J Appl Physiol* 1986; 61:2067-2073.
15. Gilman SC, Colton JS, Dutka AJ, Boogaard JS. Effect of high pressure on the release of excitatory amino acids by brain synaptosomes. *Undersea Biomed Res* 1986; 13:397-406.
16. Augustine GJ, Charlton MP, Smith SJ. Calcium action in synaptic transmitter release. *Ann Rev Neurosci* 1987; 10:633-693.
17. Huttner WB, Schiebler W, Greengard P, DeCamilli P. Synapsin I (protein I), a nerve terminal-specific phosphoprotein. III. Its association with synaptic vesicles studied in a highly purified synaptic vesicle preparation. *J Cell Biol* 1983; 96:1374-1388.
18. Liinas R, McGuinness TL, Leonard CS, Sugimori M, Greengard P. Intraterminal injection of synapsin I or calcium/calmodulin-dependent protein kinase II alters neurotransmitters release at squid giant synapse. *Proc Natl Acad Sci USA* 1985; 82:3035-3039.
19. Akerman KEO, Nicholls DG. Calcium transport by intact synaptosomes. Influence of ionophore A23187 on plasma-membrane potential, plasma membrane calcium transport, mitochondrial membrane potential, respiration, cytosolic free calcium concentration and noradrenaline release. *Eur J Biochem* 1981; 115:67-73.
20. Reed PW, Lardy HA. A23187: A divalent cation ionophore. *J Biol Chem* 1972; 247:6970-6977.
21. Neff NH, Tozer TN, Brodie BB. Application of steady-state kinetics to studies of the transfer of 5-hydroxyindoleacetic acid from brain to plasma. *J Pharmacol Exp Ther* 1967; 158:214-218.
22. Van Der Poel FW, Van Praag HM, Kork J. Evidence for a probenecid-sensitive transport system of acid monoamine metabolites from the spinal subarachnoid space. *Psychopharmacology* 1977; 52:35-40.
23. Brownstein MJ. Serotonin, histamine and the purines. In: Siegel GJ, Albers RW, Agranoff BW, Katzman R, eds. *Basic neurochemistry*, third edition. Boston: Little Brown and Co. 1981:219-231.
24. Bennett JP, Solomon SH. Serotonin synaptic receptors in the mammalian central nervous system. *Fed Proc* 1978; 37:137-138.
25. Haigler HJ, Aghajanian GK. Serotonin receptors in the brain. *Fed Proc* 1977; 36:2159-2164.